


# Study on the connectivity between the photospheric magnetic fields and their counterparts in the upper atmosphere with Hinode and IRIS

- Sunspot light bridge
- MHD waves

Ryuichi Kanoh<sup>1,2</sup>

1. The University of Tokyo
2. ISAS/JAXA





June 17, 2015      Hinode seminar @ ISAS

# Introduction

Magnetic fields are thought to play a key role in **coronal heating** problems.

Therefore, connectivity between the **photospheric magnetic fields** and **their counterparts in the upper atmosphere** is one of essential information.

The coordination of **Hinode** observation with **IRIS** observation makes it possible to understand the connectivity.






# Dynamical horizontal magnetic fields in a sunspot light bridge observed with Hinode and IRIS

Ryuichi Kanoh<sup>1,2</sup>

Toshifumi Shimizu<sup>1,2</sup>

1. The University of Tokyo


2. ISAS/JAXA



# Subject of this study

Emergence of a photospheric horizontal fields and their counterparts

(Shimizu et al. 2009)



# Remaining tasks

- Multi wavelength observation

The UV spectral data from the **IRIS** can be used with simultaneous observations by **Hinode/SOT**. Counterparts in the upper atmosphere are one of essential information in understanding how upper atmosphere responds to evolution of magnetic field at the low atmosphere.

- Temporal evolution

Although there are many observations and studies about LBs, spectro-polarimetric observations provide the evolution of magnetic fields in the cadence **longer than a few hours**, which is much longer than typical time scale of activities.




# Observation

Target: NOAA Active Region 11836



2013/08/31 16:00UT - 18:00UT  
→ Without horizontal fields

2013/09/01 16:00UT - 18:00UT  
→ Horizontal fields are emerging




| Camera        | Filter               | Observation method | Cadence |
|---------------|----------------------|--------------------|---------|
| Hinode/FG     | Ca II H              | Imaging            | 15 sec  |
| Hinode/SP     | Fe I                 | Slit scan          | 72 sec  |
| IRIS (SJI)    | Mg II k, C II, Si IV | Imaging            | 12 sec  |
| IRIS (RASTER) | Mg II k, C II, Si IV | Slit scan          | 3 sec   |

# Temporal evolution of the magnetic fields





# Configurations of the magnetic fields







$B_z$   
1300G  $\rightarrow$  100G

$B_y$   
500G  $\rightarrow$  1000G

Inclination  
150°  $\rightarrow$  100°




Magnetic fields along  
the blue/orange slices



✱ Milne-Eddington inversion is used.






# Morphology of the horizontal fields

09-01T16:00:32 Magnetic Field



Red patches

Blue patches




$\Omega$ -loop structure is identified. (If the  $180^\circ$  ambiguity is neglected)  
Flows are along horizontal fields.

Minutes after 09-01T16:00



# Properties of the up/down patches




(1)  
 $B_{\text{DOWN}} > B_{\text{UP}}$

(2)  
 $I_{\text{DOWN}} < I_{\text{UP}}$

(3)  
 $V_{\text{DOWN}} > V_{\text{UP}}$


In sunspot area, correlation between field strength and photospheric intensity is reported (Cowling 1953).

Therefore, (2) supports the result of (1).



(3) is derived by 2-component fit.


# Schematics of the photosphere




What is happening in the upper atmosphere?

$\Omega$ -shaped flux tube emerges from below the cusp-like ambient fields.  
Materials are flowed along the horizontal fields.

# Activities in the upper atmosphere




# Horizontal fields and brightening



A bright feature can be seen in the LB on September 1. Location of the feature is similar to the photospheric horizontal fields. In contrast, there are neither the bright feature nor the horizontal fields on August 31.

# Discussions[1]

## Heating mechanisms




When a horizontal flux tube emerges from below the photosphere, the interaction between ambient vertical fields and newly emerging horizontal fields may create current sheets and cause **magnetic reconnection**.



## Discussions[2]

### Driving mechanisms of the photospheric flows



These flows can be understood as a consequence of siphon flow mechanism in case of magnetohydrostatic.


✂

There is possibility that convection in inclined magnetic field geometries is a driver. However, in that case, driver that accelerates the vigorous flows is still to be identified.

# Summary

- Observed horizontal fields in a sunspot light bridge have  $\Omega$ -loop structure.
- Rapid flows are excited along the horizontal fields.
- The horizontal fields influence to heat the solar atmosphere.

# Study on dissipating processes of MHD waves with Hinode and IRIS



Ryuichi Kanoh<sup>1,2</sup>

Toshifumi Shimizu<sup>1,2</sup>


Shinsuke Imada<sup>3</sup>

1. The University of Tokyo

2. ISAS/JAXA

3. Nagoya University

# Coronal heating by means of MHD waves




There are many observational studies about MHD waves. However, study which detects **dissipating or exciting processes** of MHD waves is very rare.

Moreover, quantitative study with **spectro-polarimetric observation** will be important in the future. (In anticipation of Solar-C)


# Our idea for estimating the dissipated energy of MHD waves

(1) SOT/SP can estimate poynting flux in the photosphere by using techniques developed in previous study (Fujimura&Tsuneta (2009))

(2) With assumption of the magnetic field strength, we can use same techniques to IRIS data. (In anticipation of Solar-C)



# Review of Fujimura&Tsuneta(2009)



Solar-X:460''  
Solar-Y:10''



✂  
They averaged physical parameters of the pixels with strong CP in the ROI.

They mentioned that these small fluctuations with same frequency are products of MHD waves driven by p-mode oscillation.




We should note that there are **two possibilities** to reproduce these observed aspects

(1) P-mode oscillation excites MHD waves

(Magnetic fields are oscillated)

(2) P-mode oscillation changes the line formation height

(Actually, magnetic fields are not oscillated)




# How to distinguish ?

- Phase relation is very important.

If oscillation is caused by opacity effect, phase difference between  $B$  and  $I_c$  should be  $0^\circ$ .

In previous study, by using inverse Fourier transform **only to strong peak**, they checked phase relations and derived Poynting flux.



# Our opinion


- Phase relation is very important to distinguish between the real oscillation and the apparent oscillation (opacity effect). However, we think that the **temporal resolution in previous study (67 sec) is not enough**. We need to check that using inverse Fourier transform **only to strong peak** is reasonable or not.
- Moreover, there are some possibilities that we can obtain new aspects of MHD waves from their waveform.  
(ex. Processes of nonlinearization, attenuation etc...)

Therefore, we performed new observation on 2015/02/12 with **21 sec** cadence.

# Observations and data analysis

2015-02-12T12:25:42

Red: IRIS slit  
Cyan: SOT/SP FOV






Milne-Eddington inversion is used for deriving physical parameters.  
We averaged physical parameters of the pixels with strong CP in the ROI.



# Results



We averaged physical parameters of the pixels with strong CP in the ROI.





Subtract  
running average



- Black: Filling factor  $\times$  LOS Field strength
- Blue: Continuum Intensity
- Red: LOS velocity

# Phase relations




$I_c$  and  $B_{\text{LOS}}$  correlate positively  $\rightarrow$  consistent to opacity effect



Note that the correlation coefficient is not so large.  
There are some possibilities that influences of MHD waves are lurking.



# Comparison to the previous study



We apply Fourier transform to original profiles in the same way as previous study.

Interestingly, the results show that **only**  $B_{\text{LOS}}$  and  $v_{\text{LOS}}$  have a strong common peak. It means that  $I_c$  does not oscillate with same frequency as  $B_{\text{LOS}}$ , namely, these are **inconsistent to the opacity effect**.

# Discussions

- P-mode oscillation **does not maintain perfectly constant frequency**. Therefore, we think that focusing only to specific frequency as previous study may not be reasonable.
- It is important to distinguish the opacity effect and the real oscillation for estimating dissipated energy of the MHD waves.

# Interim summary

- Oscillating signals can be seen in  $B_{\text{LOS}}, v_{\text{LOS}}, I_{\text{C}}$ .
- These phase relations are consistent to opacity effect. This result is different from previous study.
- Focusing only to specific frequency as previous study may not be reasonable.

